

<b>Specimen No.:</b> <b>Pet/Owner:</b> <b>Service:</b> <b>Serial #:</b> <b>Expires:</b>	<b>Veterinarian:</b>
---	----------------------

Vial Sequence:      **GREEN > BLUE > RED**

**Read and follow all instructions detailed on back of page.**

For Double Treatment Sets--Sets 1 and 2 should be given concurrently.

Vial	Injection Number	Dose (cc)	Days Between Injection	Itch Level (0-10)	Date of Injection/Comments
<b>GREEN</b>  (1:1,750, wt/vol)	1	0.20	3-4		
	2	0.40	3-4		
	3	0.60	3-4		
	4	0.80	3-4		
	5	1.00	3-4		
	6	1.00	3-4		
<b>BLUE</b>  (1:175, wt/vol)	7	0.20	4-7		
	8	0.40	4-7		
	9	0.60	4-7		
	10	0.80	4-7		
	11	1.00	4-7		
	12	1.00	4-7		
<b>RED</b>  (1:35, wt/vol)	13	0.10	7-8		
	14	0.20	7-8		
	15	0.40	7-8		
	16	0.60	7-8		
	17	0.80	14-21		
	18	1.00	14-21		
	19	1.00	14-21		
	20	1.00	14-21		
	21*	1.00	14-21		
	22	1.00	14-21		

\*Contact your veterinarian to order your treatment refill at this time.

**Instructions for Administration:**

1. A weight-appropriate dose of oral antihistamine, given 45 minutes prior to each allergy injection, may increase the comfort level for the patient.
2. Clean the injection site with a moistened cloth or sterile wipe prior to the injection.
3. Using a new sterile needle and syringe, draw up the amount listed for the injection number being given. Never re-use a syringe.
4. All injections should be given subcutaneously. After insertion of the needle, withdraw plunger slightly. If blood appears in syringe, remove it and re-insert at another site.
5. Following injection, clean the injection site with a moistened cloth or sterile wipe to remove any allergens that may remain on the skin surface; this may reduce the occurrence of local inflammatory reactions.
6. Observe the animal for at least 30 minutes following each injection (see REACTIONS on back). Localized itching with redness may occur at the injection site. This is a common occurrence, and a topical, over the counter antihistamine applied on the affected skin may offer relief.
7. In the event of a severe reaction, discontinue injections and consult your veterinarian.